

IMAGES IN CLINICAL MEDICINE

Lindsey R. Baden, M.D., *Editor*

Superior Vena Cava Syndrome



Bharat Kumar, M.D.
Nael Aboul Hosn, D.O.

University of Kentucky
Lexington, KY
bharat.kumar@uky.edu

A 59-YEAR-OLD MAN WITH HYPERTENSION AND CROHN'S DISEASE THAT was complicated by fistulas was found unconscious at home. He was being treated with infliximab administered through a central venous access device, which had been placed for long-term intermittent treatment. The patient had been treated unsuccessfully with sulfasalazine, glucocorticoids, and immunomodulatory therapies, including methotrexate, azathioprine, and mercaptopurine. He was intubated and brought to the emergency department. Vital signs were stable, and the physical examination was notable for cyanosis of the head, neck, upper torso, and arms. Venography showed occlusion of the superior vena cava, a finding consistent with the superior vena cava syndrome (SVCS). Ultrasound-accelerated thrombolysis was performed along with bare-metal stenting of the superior vena cava, which alleviated the obstruction. After 2 weeks of hospitalization, the patient recovered with only residual visual impairment in the right eye (20/200) as a result of prolonged congestion of the retinal veins. SVCS is often due to external compression from cancers (e.g., cancers of the lung and lymphomas) but may be caused by intravascular thrombosis associated with the placement of a central catheter.

DOI: 10.1056/NEJMicm1311911

Copyright © 2014 Massachusetts Medical Society.